


IN THE CLAIMS:

1. (Cancelled)


2. (Currently amended) The ~~pharmaceutical agent according to Claim 1, method of claim 5 wherein the pharmaceutical agent comprises~~ a compound of formula (Ia):



~~in which the groups R^1 to R^3 have the meaning specified in Claim 1, and wherein R^9 is an alkyl group having 1-4 C atoms which, optionally, are substituted or replaced~~
by with halogen or replaced by halogen;

or ~~of~~ a pharmaceutically acceptable salt ~~of~~
such a compound thereof.


3. (Currently amended) The ~~pharmaceutical agent according to Claim 1, comprising the method of~~
claim 5 wherein the pharmaceutical agent comprises a
compound of formula (III):



or ~~of~~ a pharmaceutically acceptable salt ~~of such a~~
compound thereof.

4. (Cancelled)

5. (Currently amended) A ~~chemotherapeutic~~ method for a chemotherapeutic treatment of neuropathies characterized by application to a patient of a pharmaceutical agent comprising a compound of formula (I):



in which

R^1 =C₁₋₆alkyl, optionally substituted with halogen,

R^2 =hydrogen or C₁₋₄alkyl, optionally substituted with halogen or replaced with halogen,

R^3 =C₂₋₄alkyl, optionally substituted with halogen,

R^4 =SO₂NR⁵R⁶,

C₁₋₄alkyl, optionally substituted with NR⁵R⁶, CN, CONR⁵R⁶, CO₂R⁷, or halogen,

C₂₋₄-alkenyl, optionally substituted with NR⁵R⁶, SONR⁵R⁶, CONR⁵R⁶, CO₂R⁷, or halogen,

C₂₋₄-alkanoyl, optionally substituted with NR⁵R⁶, SONR⁵R⁶, CONR⁵R⁶, CO₂R⁷, or halogen,

R⁵ and R⁶, independent of one another, represent hydrogen or C₁₋₄alkyl, or, together with the nitrogen atom to which they are attached, represent a pyrrolidino, piperidino, morpholino, 4-(NR⁸)-1-pipera-

zinyl or 1-imidazolyl ring which, optionally, may be substituted with one or two C₁₋₄alkyl groups,

R⁷=hydrogen or C₁₋₄alkyl, optionally, substituted with fluorine, and

R⁸=hydrogen, C₁₋₃alkyl, or hydroxy alkyl having 1-4 C atoms, or ~~of~~ a pharmaceutically acceptable salt ~~of such a compound~~ thereof.

6. (Previously presented) The method of claim 5, wherein from 1-100 mg/day of said pharmaceutical agent is administered to a patient being treated.

7. (Previously presented) The method of claim 5, wherein from 5-50 mg/day of said pharmaceutical agent is administered to a patient being treated.

8. (Previously presented) The method of claim 5, wherein from 25-50 mg/day of said pharmaceutical agent is administered to a patient being treated.

9. (New) The method of claim 5 wherein the neuropathy comprises a peripheral diabetic polyneuropathy.

10. (New) The method of claim 5 wherein the neuropathy comprises gastroparesis.